TADCASTER RURAL DISTRICT COUNCIL.

ANNUAL REPORT

(INTERIM)

OF THE

Medical Officer of Health

For the Year 1943.

SELBY:
E. M. Rimmington Ltd., Printers, Booksellers and Stationers.

1944.

To:--

The Chairman and Members of the TADCASTER RURAL DISTRICT COUNCIL.

The Minister of Health has intimated that a short Interim Report only should be prepared for the year 1943.

W. BERTRAM HILL, Medical Officer of Health.

INFECTIOUS DISEASE.

The usual Table (II) is appended showing the number of Notificatione of acute Infectious Disease and the number of cases of Tuberculosis.

			Scarlet	Cereb	ro-Spinal
	Di	htheria	Fever	Erysipelas	Fever
Average for 5 year					
(1934—1938)	••••	53.6	57.6	11.6	_
1939	••••	13	43	1	_
1940	••••	17	<i>7</i> 1	3	5
1941	••••	4	47	8	4
1942	••••	8	65	4	_
1943	••••	3	40	1	

TUBERCULOSIS.

Notifications I	n		Pυ	lmonary	Non-Pulmonary
1938	••••	••••	••••	15	12
1939	••••	••••	••••	10	4
1940	••••	••••	••••	12	8
1941		••••	****	10	6
1942	••••	••••	••••	13	6
1943	••••	••••	••••	17	4

MINOR INFECTIOUS DISEASES.

Measles was epidemic during the first half of the year, the Parishes mainly involved being Tadcaster, Sherburn, South Milford, Aberford, Micklefield, Copmanthorpe, Bishopthorpe and Barwick.

69 cases of Whooping Cough were notified.

CASE RATE PER 1,000 LIVING OF ACUTE INFECTIOUS DISEASE.

	England and Tadcast										
			W	ales.		R. D.					
Typhoid		••••	••••	.01		.00					
Cerebro-Spinal Fe	ever		••••	.08		.00					
Scarlet Fever	••••	••••	••••	3.01		1.69					
Whooping Cough			••••	2.54	••••	2.93					
Diphtheria	••••	••••	••••	.88		.12					
Erysipelas		••••		.31		.04					
Small-pox		••••		.00	••••	.00					
Measles	••••	••••		9.88	••••	14.53					
Pneumonia	••••			1.34		.12					
Puerperal Pyrexia		••••		11.68	••••	2.48					

DIPHTHERIA IMMUNISATION.

It is estimated that 50% of Children under 5 and 75% above 5 years of age have been immunised.

WATER SUPPLY.

The Water Supply of the Rural District is derived as follows:—

85.5% of the houses are supplied from three sources—

- (a) The City of Leeds.
- (b) York Waterworks Company.
- (c) A Deep Bore in the Sandstone at Bilbrough owned by the District Council.

These supplies are constant, and the water is chlorinated. Bacteriological examinations are made at frequent intervals and in one source for plumbo-solvency.

The quality of the water from these three sources is satisfactory.

2.4% of the houses are supplied from privately owned sources. The water from these supplies is examined bacteriologically at frequent intervals. The quality of the water is satisfactory.

12.9% of the houses are supplied from Wells, shallow and deep. The quality of the water from these wells varies in wide degrees.

GOVERNMENT EVACUATION SCHEME.

	At the end of the year ther	e were in th	e Di	strict:-	_
(a)	Unaccompanied Children	****	****	••••	21
(b)	Mothers (17) with Children	1 (30)	••••	••••	47
(c)	Other Adults	••••	••••	••••	5
(d)	In Requisitioned Houses	Children	••••	••••	27
	Communal Billet	Adults	••••	••••	16
(e)	Other Adults	••••		••••	6

VITAL STATISTICS.

		Tadcaster	Rural trict	Eng	land and Wales.
D .1 D . 1000		1713		,	
Death Rate per 1,000	• • • •	••••	12.0		12.1
Birth Rate per 1,000	••••	••••	16.6	••••	16.5
Zymotic	••••	••••	.00		
J	••••	••••	.38		
Non-Pulmonary Tubercul			.25		
Respiratory	••••	••••	1.31		
Cancer	••••	••••	1.61		
Infantile Mortality Rate			30.6	••••	49.
(i.e., Deaths under 1 year pe	r 1,00	00 Births).			

TADCASTER RURAL DISTRICT.

Cases of Infectious Disease notified during the year 1943.

Notifiable Dispase.				Number of Cases Notified								TOVED
				es.	At all Ages—Years.							ASES REM HOSPITAL
				At all Ages.	Under 1	1 and under 5	5 and under 15	15 and under 25	25 and under 45	45 and under 65	65 and	TOTAL CASES REMOVED TO HOSPITAL
Small-pox			• •,•									
Cholera (C) Plague (P)	•••											
Diphtheria (including Mem	bran	ous Cr	oup)	3	•••	1		1	1			3
Erysipelas	•••		•	1				1			• • • •	
Scarlet Fever	•••	•••	• • •	40	•••	9	24	3	4			28
Typhus Fever	•••	•••	• • •	•••	•••		•••	•••			•••	
Enteric Fever		T3	(0)	•••		• • • •	•••	•••				•••
Relapsing Fever (R) Conti	nuea	Fever	. (C)			• • • •	• • • •	•••		•••	•••	
Puerperal Pyrexia	•••	•••	• • • •	1	•••	•••			1			1
Cerebro-spinal Meningitis	•••	•••	•••	•••	•••	•••		*****			•••	
Poliomyelitis	•••	•••	•••		2	1	••	•••			•••	
Ophthalmia Neonatorum	• • •	•••	•••	$\begin{vmatrix} 2\\17 \end{vmatrix}$		3	••				•••	1
Pulmonary Tuberculosis Other forms of Tuberculos	•••	•••	•••	4	•••	1	2	6	6	2		•••
Measles	18	•••	•••	342	15	112	119	9	6		***	•••
Primary Pneumonia	•••	•••	•••	4		112		ľ	1	1	1	1
Influenzal Pneumonia	•••	•••	•••		•••		•••	•••	_	1		1
Malaria	•••	•••	•••	•••	•••		•••	••	•••	•••	•••	•••
D	•••	•••	•••	ii	•••	4	3	•••	ï	2	1	4
Whooping Cough	•••	•••		66	5	28	31	1	ĺ			**
Trench Fever	•••	•••	•••					_		•••	•••	•••
Acute Polio Encephalitis		•••	• • • •	• • • •	•••					•••		•••
Encephalitis Lethargica			•••		•••							
•												
Tota	als	•••	•••	491	22	159	259	22	21	6	2	38